ELSEVIER

Journal of Organometallic Chemistry 637–639 (2001) 869–875

Journal of Organometallic Chemistry

www.elsevier.com/locate/jorganchem

Contents of Volumes 637–639

Foreword

R.D. Adams 1

Personal recollections

Ferrocene—how it all began
P.L. Pauson 3

How metallocene chemistry and research began in Munich
E.O. Fischer, R. Jira 7

The early ferrocene days—a personal recollection
M. Rosenblum 13

Recollections of the arrival of ferrocene
M.C. Whiting 16

Cyclopentadienyl–metal chemistry in the Wilkinson Group, Harvard, 1952–1955
F.A. Cotton 18

Reviews

Studies on the cyclometallation of ferrocenylimines
Y. Wu, S. Huo, J. Gong, X. Cui, L. Ding, K. Ding, C. Du, Y. Liu, M. Song 27

Ferrocene polymers: current architectures, syntheses and utility
R.D.A. Hudson 47

Communications

The reaction of ferrocenyl acetylene with $[RuCp(PR_3)(CH_3CN)_2]PF_6$ (R = Me, Ph, Cy). Formation of the first allenyl carbene complexes
E. Rüba, K. Mereiter, R. Schmid, K. Kirchner, H. Schottenberger 70

Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy)
T. Moriuchi, K. Yoshida, T. Hirao 75

Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication
Y. Yamada, J. Mizutani, M. Kurihara, H. Nishihara 80

[2 + 2 + 2] Cyclotrimerisation of bisaryl acetylene bearing ferrocenyl units with planar chirality: synthesis of enantiopure conjugated polyferrocene complexes
V. Mamane, A. Gref, F. Lefloch, O. Riant 84

Metal-directed assembly of polyferrocenyl transition metal dithiocarbamate macrocyclic molecular boxes
S.-W. Lai, M.G.B. Drew, P.D. Beer 89

Regular papers

Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety
H. Runqiu, W. Qingmin 94

ortho-Silylation of 2,2′-bis(oxazolinyl)-1,1′-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation
S. Lee, J.H. Koh, J. Park 99

Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents
P.A. Deck, C.E. Kroll, W. Gary Hollis Jr., F.R. Fronczek 107

Chiral ferrocenylthiazolidines, new ligands for palladium complexes
A. González, J.R. Granell, C. López 116

1,1′,2,2′-Tetrakis(diphenylphosphino)-4,4′-di-*tert*-butylferrocene, a new *cisoid* arrangement of phosphino groups
R. Broussier, E. Bentabet, R. Amardeil, P. Richard, P. Meunier, P. Kalck, B. Gautheron 126

Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents
O.B. Sutcliffe, A. Chesney, M.R. Bryce 134

Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry
K.R. Justin Thomas, J.T. Lin 139

Ruthenium(II) complexes containing 4-ferrocenylphenylisocyanide ligands. Crystal structure of *trans*, *trans*, *trans*-$[RuCl_2(POMe\text{–}P)_2(FcC_6H_4NC)_2]$ ($POMe = PPh_2C_6H_4OCH_3$; Fc = ferrocenyl)
O. Clot, M.O. Wolf, G.P.A. Yap 145

An electrochemical study in acetonitrile of macrocyclic or open-chain ferrocene-containing oxa-aza or polyaza receptors in the presence of protons, metal cations and anions
J.M. Lloris, R. Martínez-Máñez, J. Soto, T. Pardo. 151

Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers
W.-Y. Wong, G.-L. Lu, K.-F. Ng, C.-K. Wong, K.-H. Choi 159

Iron(II) versus osmium(II) oxidation in 1,1′-bis(diorganophosphino)ferrocene–osmium(II) complexes
W. Kaim, T. Sixt, M. Weber, J. Fiedler 167

Bonding in 'closed,' open, and half-open ferrocenes: new insight from structural and Mössbauer spectroscopic studies
R. Basta, D.R. Wilson, H. Ma, A.M. Arif, R.H. Herber, R.D. Ernst 172

(Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties
B. Malézieux, R. Andrés, M. Brissard, M. Gruselle, C. Train, P. Herson, L.L. Troitskaya, V.I. Sokolov, S.T. Ovseenko, T.V. Demeschik, N.S. Ovanesyan, I.A. Mamed'yarova. 182

Facile synthesis of bidimensional ferrocenyl-based branched oligomers by palladium-catalyzed coupling reactions
A. Peruga, J.A. Mata, D. Sainz, E. Peris 191

4-Ethinyl-benzonitrile-ferrocenes bridged by a $Pd(PPh_3)_2$ unit; the solid-state structure of $(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_4C{\equiv}CC_6H_4C{\equiv}N\text{-}1,4)$
S. Köcher, H. Lang 198

Synthesis and molecular structure of a novel ferrocene-containing macrocyclic acyl thiourea derivative
L.-y. Zhang, Y.-f. Yuan, A.-g. Hu, J.-t. Wang, J. Sun 204

Synthesis, structures and electrochemical properties of Group 6 metal carbonyl complexes containing ferrocenylpyrazole ligands
L.-F. Tang, W.-L. Jia, Z.-H. Wang, J.-F. Chai, J.-T. Wang. 209

Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids
S.R. Alley, W. Henderson 216

Synthesis of controlled π-extended conjugate nanostructures of 1,1′-ferrocene
J.-G. Rodríguez, S. Pleite. 230

Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit
S. Hartmann, R.F. Winter, T. Scheiring, M. Wanner 240

Ferrocene as a ferromagnetic coupler. Synthesis and characterization of a ferrocene bridged polychlorotriphenylmethyl diradical
O. Elsner, D. Ruiz-Molina, I. Ratera, J. Vidal-Gancedo, C. Rovira, J. Veciana 251

Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α-bromoacetyl substituted ferrocenes
P. Molina, A. Tárraga, D. Curiel, M.D. Velasco 258

An investigation of the chemistry of ferrocenoyl derivatives. The synthesis and reactions of ferrocenoyl imidazolide and its derivatives
C. Imrie, L. Cook, D.C. Levendis 266

Synthesis, characterisation, crystal structures and electrochemistry of triosmium nitrite carbonyl clusters containing ferrocenyl phosphines
E.N.-M. Ho, B.K.-M. Hui, W.-T. Wong 276

Synthesis and characterization of ferrocenylalcohol derivatives of hexachlorocyclotriphosphazene. X-ray crystal structure of $N_3P_3Cl_5OCH_2CH_2C_5H_4FeCp$
C. Nataro, C.N. Myer, W.M. Cleaver, C.W. Allen 284

Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study
P. Štěpnička, L. Trojan, J. Kubišta, J. Ludvík 291

Synthesis and structure of an alkylgallium bridged [1,1]ferrocenophane
W. Uhl, I. Hahn, A. Jantschak, T. Spies 300

The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands
J.D. Carr, S.J. Coles, M.B. Hursthouse, J.H.R. Tucker 304

Electrostatic field effects manifested in ferrocenyl metal complexes and the crystal structure of $[Fe(\eta^5\text{-}C_5H_5)(\eta^5\text{-}C_5H_4CH{=}NNHC_5H_4N)]\cdot HCl$
J. Silver, G.R. Fern, J.R. Miller, E. Slade, M. Ahmet, A. Houlton, D.J. Evans, G.J. Leigh 311

Study on the synthesis of nonracemic C_2-symmetric 1,1′-binaphthyl-2,2′-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy
P. Kasák, R. Mikláš, M. Putala 318

Properties of some ferrocene macrocyclic dioxotetraamines: the roles of aromatic side-arms
P. Xue, Q. Yuan, E. Fu, C. Wu 327

1,1′-Ferrocenoyl–oligoprolines. A synthetic, structural and electrochemical study
Y. Xu, P. Saweczko, H.-B. Kraatz 335

Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes
J.L. Sessler, R.S. Zimmerman, G.J. Kirkovits, A. Gebauer, M. Scherer 343

Solution phase peptide synthesis with ferrocenyl amino acid derivatives
J. Sehnert, A. Hess, N. Metzler-Nolte 349

Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands
O. Reynes, F. Maillard, J.-C. Moutet, G. Royal, E. Saint-Aman, G. Stanciu, J.-P. Dutasta, I. Gosse, J.-C. Mulatier 356

Highly efficient reduction of ferrocenyl derivatives by borane
L. Routaboul, J. Chiffre, G.G.A. Balavoine, J.-C. Daran, E. Manoury 364

Ferrocenes derived from cyclopenta[*l*]phenanthrene: dibenzindene–metal complexes that resist haptotropic shifts
S.S. Rigby, A. Decken, A.D. Bain, M.J. McGlinchey 372

Reactions of ferrocene-derived bis(hydroxymethyl) phosphine sulfides $FcCH(R)P(S)(CH_2OH)_2$ ($R = H$, CH_3) with cyclic thionylphosphazenes: crystal structures of $FcCH_2P(S)(CH_2O)_2PN(NPCl_2)[NS(O)Ph]$ and $FcCH_2P(S)(CH_2O)_2PN_2P[N(Me)CH_2]_2[NS(O)Ph]$ (Fc = ferrocenyl)
T.V.V. Ramakrishna, A.J. Elias 382

Multistep redox properties of 2,2′-bipyridylboronium substituted ferrocenes
L. Ding, K. Ma, M. Bolte, F. Fabrizi de Biani, P. Zanello, M. Wagner 390

Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides
A.E. Beilstein, M.W. Grinstaff 398

Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes
B.F. Bonini, M. Comes-Franchini, M. Fochi, G. Mazzanti, A. Ricci, M. Tomasulo, G. Varchi 407

Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes
K. Bushell, C. Gialou, C.H. Goh, N.J. Long, J. Martin, A.J.P. White, C.K. Williams, D.J. Williams, M. Fontani, P. Zanello 418

Knoevenagel condensation of ferrocenyl substituted carboxaldehydes with methylene active compounds on inorganic supports and the NLO properties of the resulting push–pull alkenes
E. Stankovic, S. Toma, R. Van Boxel, I. Asselberghs, A. Persoons 426

The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, $[Fe_2(\eta^5\text{-}C_5H_5)_2(CO)_2(\mu\text{-}CO)(\mu\text{-}C\text{–}CH_3)]^+[BF_4]^-$ and terthienyl spacers. Crystal structure of 2-[(*E*)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene
R.D.A. Hudson, I. Asselberghs, K. Clays, L.P. Cuffe, J.F. Gallagher, A.R. Manning, A. Persoons, K. Wostyn 435

Electron acceptors of the fluorene series: Part 12. 9-(Metalloceneylidene)nitrofluorene derivatives of Fc–NF, NF–Fc–NF, and NF–Rc–NF types, and the vinylogues Fc–π–NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene–ferrocene derivatives
D.F. Perepichka, I.F. Perepichka, A.F. Popov, M.R. Bryce, A.S. Batsanov, A. Chesney, J.A.K. Howard, N.I. Sokolov. 445

Octahedral ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(S_p)-2-(diphenylphosphino)ferrocenyl]-4-(isopropyl)oxazoline (FcPN). X-Ray crystal structures of *fac*-$[RuCl_2(PMe_3)_2(FcPN)]$ and *fac*-$[RuCl_2(dppm)(FcPN)]$ (dppm = bis(diphenylphosphino)methane)
J. Gimeno, E. Lastra, C. Madrigal, C. Graiff, A. Tiripicchio 463

Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations
A.D. Ryabov, E.S. Ryabova, M.D. Reshetova 469

Ferrocenyl naphthalene diimide can bind to DNA·RNA hetero duplex: potential use in an electrochemical detection of mRNA expression
S. Sato, S. Fujii, K. Yamashita, M. Takagi, H. Kondo, S. Takenaka 476

Syntheses and some reactions of complexes containing carbon chains capped by ferrocenyl and $W(CO)_3Cp$ groups
M.I. Bruce, M.E. Smith, B.W. Skelton, A.H. White 484

Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series
S. Top, A. Vessières, C. Cabestaing, I. Laios, G. Leclercq, C. Provot, G. Jaouen 500

Ferrocene substituted nitronyl nitroxide and imino nitroxide radicals. Synthesis, X-ray structure and magnetic properties
C. Sporer, D. Ruiz-Molina, K. Wurst, H. Kopacka, J. Veciana, P. Jaitner 507

A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with $Os_3(CO)_{10}(\mu\text{-py})(\mu\text{-H})$
R.D. Adams, B. Qu, M.D. Smith 514

Synthesis, reactivity, and crystal structures of ferrocene-substituted amidinate derivatives
J.R. Hagadorn, J. Arnold 521

New chiral ferrocenyl-pyridinium salts for non-linear optics
K. Roque, F. Barangé, G.G.A. Balavoine, J.-C. Daran, P.G. Lacroix, E. Manoury. 531

Diferrocenyltriphosphines 2. Reversible phosphine deligation in the chemistry of diferrocenyltriphosphine Ru(II) dichloride complexes with nitriles and pyridines: towards a pH-switchable catalyst?
I.R. Butler, S.J. Coles, M. Fontani, M.B. Hursthouse, E. Lewis, K.L.M.A. Malik, M. Meunier, P. Zanello. 538

$(\eta^5\text{-}C_5H_5)$-Ring alkylation reaction with the C_5H_5 anion: towards the construction of tri-Fe complex $Fe\{[\mu,\eta^5\text{:}\eta^4\text{-}5\text{-}exo\text{-}(1'\text{-}C_5H_4)C_5H_5]Fe(CO)_2(PPh_3)\}_2$
L.-S. Luh, L.-K. Liu 549

Semimasked 1,1′-diethynylferrocenes: synthetic concepts, preparations, and reactions
H. Schottenberger, J. Lukassser, E. Reichel, A.G. Müller, G. Steiner, H. Kopacka, K. Wurst, K.H. Ongania, K. Kirchner 558

Novel cyclopalladated ferrocenyl Schiff base compounds with bridging and chelating diphosphines. Crystal and molecular structure of $[\{Pd[(\eta^5\text{-}C_5H_5)Fe(\eta^5\text{-}C_5H_3)C(H)\text{=}N\text{-}2,4,6\text{-}Me_3C_6H_2]\}\{Ph_2P(CH_2)_nPPh_2\text{–}P,P\}][PF_6]$ ($n = 1, 2$)
J.M. Vila, E. Gayoso, T. Pereira, M. Mariño, J. Martínez, J.J. Fernández, A. Fernández, M. López-Torres 577

A re-examination of the reaction between 1,1′-bis(diphenylphosphino)ferrocene (dppf) diselenide and $[Ru_3(CO)_{12}]$. Electrochemical behaviour of the isomeric *nido*-clusters $[Ru_3(\mu_3\text{-}Se)_2(dppf)(CO)_7]$ and $[Ru_3(\mu_3\text{-}Se)_2(\mu\text{-}dppf)(CO)_7]$ and crystal structure of $[Ru_3Se\{\mu\text{-}P(Ph)C_5H_4FeC_5H_4PPh_2\}(\mu\text{-}OCPh)(CO)_6]$
F.F. de Biani, C. Graiff, G. Opromolla, G. Predieri, A. Tiripicchio, P. Zanello 586

Water soluble ferrocenyl and polyferrocenyl compounds: synthesis and electrochemistry
A. Salmon, P. Jutzi 595

Organometallic building blocks for crystal engineering. Synthesis, structure and hydrogen bonding interactions in [Fe(η^5-C_5H_4–$CH_2(CH_3)OH)_2$], [Fe(η^5-$C_5H_3(CH_3)COOH)_2$], [Fe(η^5-$C_5H_4CH(CH_3)NH(\eta^5$-$C_5H_4CH(CH_3)$)] and in the diaminecyclohexane salt $[Fe(\eta^5\text{-}C_5H_4COO)_2]^{2-}[(1S,2S)\text{-}(NH_3)_2C_6H_{10}]^{2+}\cdot 2[H_2O]$
D. Braga, L. Maini, F. Paganelli, E. Tagliavini, S. Casolari, F. Grepioni 609

Ferrocenyl substituted chlorostilbenes and butadienes
K. Senthil Kumar, K.C. Kumara Swamy 616

Formation and structural characterization of amino-substituted [3]ferrocenophanes derived from intramolecular Mannich-type coupling reactions at the metallocene framework
P. Liptau, S. Knüppel, G. Kehr, O. Kataeva, R. Fröhlich, G. Erker 621

Preparation, characterization, electrochemical studies and crystal structure determination of salicylaldehyde–aroylhydrazone, ferrocenyl–aroylhydrazone and salicylaldehyde–ferrocenoylhydrazone complexes of indium
Y. Yuan, Z. Cao, N. Fu, J. Wang, L. Weng, A. Bezerra de Carvalho, C. Peppe 631

Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units
B. Alonso, B. González, B. García, E. Ramírez-Oliva, M. Zamora, C.M. Casado, I. Cuadrado 642

Bis(ferrocenyl)mercury as a source of ferrocenyl moiety in Pd-catalyzed reactions of carbon–carbon bond formation
I.P. Beletskaya, A.V. Tsvetkov, G.V. Latyshev, V.A. Tafeenko, N.V. Lukashev 653

Synthesis and characterization of new germylferrocenyl, germylferrocenophane and polymeric germyleneferrocenylene systems
M. Castruita, F. Cervantes-Lee, J.S. Mahmoud, Y. Zhang, K.H. Pannell 664

A rapid approach to ferrocenophanes via ring-closing metathesis
A.J. Locke, C. Jones, C.J. Richards 669

Copper(I) complexes containing 1,1′-bis(diphenylphosphino) ferrocene (dppf) as a chelate ligand. X-ray crystal structures of [$Cu_2(\mu$-$Cl)_2(\kappa^2$-*P*,*P*-dppf)$_2$] and [$Cu(CN^tBu)_2(\kappa^2$-*P*,*P*-dppf)][BF_4]
J. Díez, M.P. Gamasa, J. Gimeno, M. Lanfranchi, A. Tiripicchio 677

Synthetic, electrochemical, and structural studies on heterobimetallic crown thioether complexes with Group 10 metals: the crystal structures of [Pt(9S3)(dppf)]$(PF_6)_2\cdot CH_3NO_2$ and [Pd(9S3)(dppf)]$(PF_6)_2\cdot CH_3NO_2$
G.J. Grant, S.M. Carter, A. LeBron Russell, I.M. Poullaos, D.G. VanDerveer 683

Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX_2 (M = Pt, Pd; X = Cl, Br, I)
T.J. Colacot, H. Qian, R. Cea-Olivares, S. Hernandez-Ortega 691

Asymmetric synthesis of planar chiral 2-mono- and 2,2′-disubstituted 1,1′-bisbenzoylferrocenes
D. Enders, T. Klumpen 698

Synthesis and characterisation of rhodium(0) and rhodium(−1) complexes stabilised by 1,1′-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of [Rh(dppf)$_2$] and [Na(THF)$_5$][Rh(dppf)$_2$]·THF
B. Longato, R. Coppo, G. Pilloni, C. Corvaja, A. Toffoletti, G. Bandoli 710

Notes

Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition
W.-y. Liu, Q.-h. Xu, Y.-m. Liang, B.-h. Chen, W.-m. Liu, Y.-x. Ma 719

Synthesis of 1′,1‴-bis(heterocyclyl)biferrocene compounds
C.-M. Liu, Y.-m. Liang, X.-l. Wu, W.-m. Liu, Y.-x. Ma 723

New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines
B. Wrackmeyer, H.E. Maisel, M. Herberhold 727

Synthesis and crystal structure of 1,1′-bis[4-(2,2′:6′,2″-terpyridin-4′-yl)phenyl]octamethylferrocene
U. Siemeling, U. Vorfeld, B. Neumann, H.-G. Stammler, M. Fontani, P. Zanello 733

Ferrocene derivatives (I). Synthesis and X-ray structure determination of *N*-*o*-methylphenylferrocenesulfonamide
M. Li, Y. Bai, J. Lu, B. Yang, K. Zhu, H. Ma 738

Ferrocene derivatives (II). Synthesis and reactions of 4-ferrocenyl-2-thiazolamine
H. Ma, Y. Hou, Y. Bai, J. Lu, B. Yang 742

Bis[bis(trimethylsilyl)cyclopentadienyl]selane as a ligand: selena-2,2′,4,4′-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triselena-2,2′,4,4′-tetrakis(trimethylsilyl)-[3]ferrocenophane
G. Thaler, B. Klotz, K. Wurst, F. Sladky 745

A new phthalocyanine–ferrocene conjugated dyad
A. González-Cabello, P. Vázquez, T. Torres 751

Palladium(II) phosphine thiocarboxylates. Structures of *cis*-Pd(κ*S*-SOCMe)$_2$(dppf) [dppf = 1,1′-bis(diphenylphosphino)ferrocene] and *trans*-Pd(κ*S*-SOCMe)$_2$(PPh$_3$)$_2$
Y.C. Neo, J.J. Vittal, T.S.A. Hor 757

Novel ferrocene modified *P*,*N*-ligands for enantioselective palladium-catalyzed allylic substitution reactions
S.-L. You, X.-L. Hou, L.-X. Dai 762

Reactivity of the bis(allyl)-ruthenium(IV) dimer [{Ru(η^3:η^3-C$_{10}$H$_{16}$)(μ-Cl)Cl}$_2$] (C$_{10}$H$_{16}$ = 2,7-dimethylocta-2,6-diene-1,8-diyl) towards dppf (dppf = [Fe(η^5-C$_5$H$_4$PPh$_2$)$_2$]) and dppfO (dppfO = [Fe(η^5-C$_5$H$_4$PPh$_2$)(η^5-C$_5$H$_4$P(=O)Ph$_2$)])
V. Cadierno, S.E. García-Garrido, J. Gimeno 767

Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrocenyl fragment
F. Tisato, F. Refosco, M. Porchia, G. Bandoli, G. Pilloni, L. Uccelli, A. Boschi, A. Duatti 772

Charge transfer spectra and photoreactivity of (*E*)-1-ferrocenyl-2-(1-methyl-4-pyridiniumyl)ethylene cation
H. Kunkely, A. Vogler 777

Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1′-diacetylferrocene with aromatic aldehydes in dry conditions
W.-y. Liu, Q.-h. Xu, B.-h. Chen, Y.-m. Liang, Y.-x. Ma, W.-m. Liu 782

Hindering the formation of ferrocenes: mono(cyclopentadienyl)halo iron complexes [Fe(C$_5$R$_5$)X] containing a sterically bulky cyclopentadienyl ligand
J. Okuda 786

Cyclopalladated ferrocenylimines: highly active catalysts for Heck reactions
Y. Wu, J. Hou, H. Yun, X. Cui, R. Yuan 793

The reaction of ferrocene with mercuric chloride revisited: '$Cp_2Fe*2HgCl_2$' = [(C$_5$H$_5$)Fe(C$_5$H$_4$HgCl)]$_2$[Hg$_2$Cl$_6$], comparison of its molecular structure with that of chloromercuriferrocene
K. Sünkel, T. Kießling 796

Redox behavior of boronato-functionalized 1,1′-bis(diphenylphosphino)ferrocenes
P. Zanello, A. Cinquantini, M. Fontani, M. Giardiello, G. Giorgi, C.R. Landis, B.F.M. Kimmich 800

Palladium(II) complexes containing symmetrical diphosphine ligands based on ferrocene. Sterically enforced palladium–iron bonds
M.A. Zuideveld, B.H.G. Swennenhuis, P.C.J. Kamer, P.W.N.M. van Leeuwen 805

Electrochemical determination of diffusion coefficient of π-conjugated polymers containing ferrocene unit
T. Morikita, T. Yamamoto 809

Synthesis of ferrocenylketyl radicals by chromium(II) complexes
Z.R. Ratkovic, L. Somsák, K. Micskei, C. Zucchi, G. Pályi 813

Control of the symmetry of binuclear ferrocene derivative by using chiral substituent and its mixed-valence state
T. Oda, S. Nakashima, T. Okuda 820

Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu$_4$][B(C$_6$F$_5$)$_4$] as supporting electrolyte
N. Camire, U.T. Mueller-Westerhoff, W.E. Geiger 823

Second harmonic generation in ferrocene based hydrogen bonded assemblies
S.K. Pal, A. Krishnan, P.K. Das, A.G. Samuelson 827

Short syntheses of 8-substituted 8′-[1-(1′-phenylthio)ferrocenyl]-1,1′-binaphthyls from Suzuki coupling reactions. A strategy for generating new chiral ligands and charge-transfer complexes
D.H. Hua, J.W. McGill, M. Ueda, H.A. Stephany 832

Synthesis of new ferrocene containing diamines and their use in epoxy resins
M.E. Wright, J. Laub, P.R. Stafford, W.P. Norris 837

Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions
T. Kondo, M. Okamura, K. Uosaki 841

Palladium-catalyzed asymmetric allylic alkylations of cycloalkenyl acetates with planar chiral phosphino-ferrocene carboxylic acids
S.-L. You, Y.-M. Luo, W.-P. Deng, X.-L. Hou, L.-X. Dai 845

Author index of Vol. 637–639. 851
Subject index of Vol. 637–639. 855
Contents of Vol. 637–639 869